




3. Given your unnatural passion for simultaneous equilibria, you have been assigned the task of teaching a Quantitative Analysis class about the role of charge and mass balance in equilibrium systems. Briefly define and illustrate each term, using a solution containing 0.0010 M HCl and 0.10 M oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$, $\text{pK}_{\text{a}1} = 1.252$, $\text{pK}_{\text{a}2} = 4.266$) as an example. Your answer should include example mass and charge balance expressions.

Do problem 4. (15 pts)

4. When ammonium sulfate dissolves, both the anion and the cation can participate in acid-base equilibria. Considering the equilibria below, write enough equations so that you could solve for the concentration of each species in solution. You must write the charge balance expression and at least one mass balance. A numerical result isn't necessary.




$$K_{\text{sp}} = 276$$

$$K_{\text{a}} = 5.7 \times 10^{-10}$$

$$K_{\text{b}} = 9.8 \times 10^{-13}$$

7. Clearly describe the case when it is preferable to use calibration by standard additions, rather than a traditional calibration curve for an analysis. Include an example of how you would run the experiment and extract an unknown concentration from your data.

8. Is it possible to perform a 99.9 % complete separation of barium and silver by precipitation with sulfate if both Ba^{2+} and Ag^+ are present initially at 0.020 M? Justify your decision.



Possibly Useful Information

$K_a K_b = K_w = 1.0 \times 10^{-14}$	Don't eat the yellow snow.
$-\log \gamma = \frac{0.51z^2 \sqrt{\mu}}{1 + \alpha \sqrt{\mu}/305}$	$\mu = \frac{1}{2} \sum c_i z_i^2$
$\Delta G = \Delta H - T\Delta S = -RT \ln K$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
$\frac{I_x}{I_{s+x}} = \frac{k[x]_i}{k([s]_f + [x]_f)} = \frac{[x]_i}{[s]_f + [x]_f}$	$\frac{\text{Analyte Signal}}{\text{Analyte Concentration}} = F \left(\frac{\text{Standard Signal}}{\text{Standard Concentration}} \right)$

Activity coefficients for aqueous solutions at 25°C

Ion	Ion size (α , pm)	Ionic strength (μ , M)				
		0.001	0.005	0.01	0.05	0.1
CHARGE = ± 1						
H ⁺	900	0.967	0.933	0.914	0.86	0.83
(C ₆ H ₅) ₂ CHCO ₂ ⁻ , (C ₃ H ₇) ₄ N ⁺	800	0.966	0.931	0.912	0.85	0.82
(O ₂ N) ₃ C ₆ H ₂ O ⁻ , (C ₃ H ₇) ₃ NH ⁺ , CH ₃ OC ₆ H ₄ CO ₂ ⁻	700	0.965	0.930	0.909	0.845	0.81
Li ⁺ , C ₆ H ₅ CO ₂ ⁻ , HOC ₆ H ₄ CO ₂ ⁻ , ClC ₆ H ₄ CO ₂ ⁻ , C ₆ H ₅ CH ₂ CO ₂ ⁻ , CH ₂ =CHCH ₂ CO ₂ ⁻ , (CH ₃) ₂ CHCH ₂ CO ₂ ⁻ , (CH ₃ CH ₂) ₄ N ⁺ , (C ₃ H ₇) ₂ NH ₂ ⁺	600	0.965	0.929	0.907	0.835	0.80
Cl ₂ CHCO ₂ ⁻ , Cl ₃ CCO ₂ ⁻ , (CH ₃ CH ₂) ₃ NH ⁺ , (C ₃ H ₇)NH ₃ ⁺	500	0.964	0.928	0.904	0.83	0.79
Na ⁺ , CdCl ⁺ , ClO ₂ ⁻ , IO ₃ ⁻ , HCO ₃ ⁻ , H ₂ PO ₄ ⁻ , HSO ₃ ⁻ , H ₂ AsO ₄ ⁻ , Co(NH ₃) ₄ (NO ₂) ₂ ⁺ , CH ₃ CO ₂ ⁻ , ClCH ₂ CO ₂ ⁻ , (CH ₃) ₄ N ⁺ , (CH ₃ CH ₂) ₂ NH ₂ ⁺ , H ₂ NCH ₂ CO ₂ ⁻	450	0.964	0.928	0.902	0.82	0.775
⁺ H ₃ NCH ₂ CO ₂ H, (CH ₃) ₃ NH ⁺ , CH ₃ CH ₂ NH ₃ ⁺	400	0.964	0.927	0.901	0.815	0.77
OH ⁻ , F ⁻ , SCN ⁻ , OCN ⁻ , HS ⁻ , ClO ₃ ⁻ , ClO ₄ ⁻ , BrO ₃ ⁻ , IO ₄ ⁻ , MnO ₄ ⁻ , HCO ₂ ⁻ , H ₂ citrate ⁻ , CH ₃ NH ₃ ⁺ , (CH ₃) ₂ NH ₂ ⁺	350	0.964	0.926	0.900	0.81	0.76
K ⁺ , Cl ⁻ , Br ⁻ , I ⁻ , CN ⁻ , NO ₂ ⁻ , NO ₃ ⁻	300	0.964	0.925	0.899	0.805	0.755
Rb ⁺ , Cs ⁺ , NH ₄ ⁺ , TI ⁺ , Ag ⁺	250	0.964	0.924	0.898	0.80	0.75
CHARGE = ± 2						
Mg ²⁺ , Be ²⁺	800	0.872	0.755	0.69	0.52	0.45
CH ₂ (CH ₂ CH ₂ CO ₂ ⁻) ₂ , (CH ₂ CH ₂ CH ₂ CO ₂ ⁻) ₂	700	0.872	0.755	0.685	0.50	0.425
Ca ²⁺ , Cu ²⁺ , Zn ²⁺ , Sn ²⁺ , Mn ²⁺ , Fe ²⁺ , Ni ²⁺ , Co ²⁺ , C ₆ H ₄ (CO ₂ ⁻) ₂ , H ₂ C(CH ₂ CO ₂ ⁻) ₂ , (CH ₂ CH ₂ CO ₂ ⁻) ₂	600	0.870	0.749	0.675	0.485	0.405
Sr ²⁺ , Ba ²⁺ , Cd ²⁺ , Hg ²⁺ , S ²⁻ , S ₂ O ₄ ²⁻ , WO ₄ ²⁻ , H ₂ C(CO ₂ ⁻) ₂ , (CH ₂ CO ₂ ⁻) ₂ , (CHOHCO ₂ ⁻) ₂	500	0.868	0.744	0.67	0.465	0.38
Pb ²⁺ , CO ₃ ²⁻ , SO ₃ ²⁻ , MoO ₄ ²⁻ , Co(NH ₃) ₅ Cl ²⁺ , Fe(CN) ₅ NO ²⁻ , C ₂ O ₄ ²⁻ , Hcitrate ²⁻	450	0.867	0.742	0.665	0.455	0.37
Hg ₂ ²⁺ , SO ₄ ²⁻ , S ₂ O ₃ ²⁻ , S ₂ O ₆ ²⁻ , S ₂ O ₈ ²⁻ , SeO ₄ ²⁻ , CrO ₄ ²⁻ , HPO ₄ ²⁻	400	0.867	0.740	0.660	0.445	0.355
CHARGE = ± 3						
Al ³⁺ , Fe ³⁺ , Cr ³⁺ , Sc ³⁺ , Y ³⁺ , In ³⁺ , lanthanides ^a	900	0.738	0.54	0.445	0.245	0.18
citrate ³⁻	500	0.728	0.51	0.405	0.18	0.115
PO ₄ ³⁻ , Fe(CN) ₆ ³⁻ , Cr(NH ₃) ₆ ³⁺ , Co(NH ₃) ₆ ³⁺ , Co(NH ₃) ₅ H ₂ O ³⁺	400	0.725	0.505	0.395	0.16	0.095
CHARGE = ± 4						
Th ⁴⁺ , Zr ⁴⁺ , Ce ⁴⁺ , Sn ⁴⁺	1100	0.588	0.35	0.255	0.10	0.065
Fe(CN) ₆ ⁴⁻	500	0.57	0.31	0.20	0.048	0.021

a. Lanthanides are elements 57–71 in the periodic table. SOURCE: J. Kielland, *J. Am. Chem. Soc.* **1937**, *59*, 1675.